

RAMIK® Green

(EPA REG. NO. 2393-498)

SUPPLEMENTAL LABELING**FOR USE ONLY IN THE STATE OF MONTANA****RODENTICIDE FOR CONTROL OF GROUND SQUIRRELS****ACTIVE INGREDIENT:**

Diphacinone (2-Diphenylacetyl-1,3-

Indandione 0.005%

INERT INGREDIENTS:.....99.995%

Total.....100.000%

EPA SLN No MT-860003

EPA REG No 2393-498

EPA EST No 61282-WI-1

CAUTION**KEEP OUT OF THE REACH OF CHILDREN**

Keep out of reach of children, pets, wildlife and domestic animals.

RAMIK bait should be placed in tamper-proof bait boxes or in areas which are inaccessible to children, pets, wildlife, and domestic animals.

Keep out of lakes, streams or ponds.

Wear rubber gloves when handling this product.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K₁, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

NOTICE

Because Manufacturer and Seller have no control over storage, handling, and conditions of use, which are of critical importance, Manufacturer and Seller make no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate government agencies, due to misuse, improper handling or storage of this material. Nor do Manufacturer and Seller assume any responsibility for injury to persons, crops, animals, soil,

or property arising out of misuse, improper handling, or storage of this material.

DIRECTIONS FOR USE

It is a violation of State and Federal law to use this product in a manner inconsistent with its labeling.

GROUND SQUIRRELS: (Columbian ground squirrel, *Spermophilus columbianus*) and (Richardson's ground squirrel, *S. richardsoni*)

This product can be used to control ground squirrels found in levee and ditch banks, around farm buildings, along fence lines, in orchards, in crops, or in non-crop areas.

All outdoor bait placements must be made in protected bait stations, where they will be readily accessible to the ground squirrel, preferably near their burrows.

These bait stations must be strong enough to inhibit destruction by livestock and nontarget animals. Stations must be constructed to permit entry by target species but not by larger animals. Stations must be constructed to inhibit spillage of bait to the outside environment and must be secured so that they cannot be overturned by wind, livestock, pets and wildlife.

Bait placements should be checked frequently to insure that an uninterrupted supply of bait is always available. Maintain an uninterrupted supply of bait for at least 30 days or until all signs of feeding have stopped.

Bait stations may be maintained on a permanent basis to prevent reinfestation of the area.

Bait stations should be located conveniently to the squirrel burrows and should be placed 20 to 100 feet apart, depending on population density.

Use 2 to 4 pounds of bait in each protected bait station.

STORAGE AND DISPOSAL

Please refer to container label for appropriate statements.

24(c) Registrant:

HACO, Inc. • PO Box 7190 • Madison, WI 53707